

AD-A162 957

A MARTINGALE CHARACTERIZATION OF MIXED POISSON  
PROCESSES(U) NORTH CAROLINA UNIV AT CHAPEL HILL CENTER  
FOR STOCHASTIC PROCESSES D PFEIFER OCT 85 TR-119

1/1


UNCLASSIFIED

AFOSR-TR-85-1177 F49620-85-C-0144

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

AD-A162 957

2

SECURITY CLASSIFICATION OF THIS PAGE



## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT <b>unlimited</b> Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) <b>Technical Report No. 119</b>		5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR- 85-1177</b>	
6a. NAME OF PERFORMING ORGANIZATION <b>Center for Stochastic Processes</b>	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION <b>Air Force Office of Scientific Research</b>	
6c. ADDRESS (City, State and ZIP Code) <b>Statistics Dept., Univ. of North Carolina, Phillips Hall 039-A Chapel Hill, NC 27514</b>		7b. ADDRESS (City, State and ZIP Code) <b>Bolling Air Force Base Washington, DC 20332</b>	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION <b>AFOSR</b>	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER <b>F49620 85 C 0144</b>	
8c. ADDRESS (City, State and ZIP Code) <b>Bolling Air Force Base Washington, DC 20332</b>		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. <b>61103F</b>	PROJECT NO. <b>2304</b>
		TASK NO. <b>A5</b>	WORK UNIT NO.
11. TITLE (Include Security Classification) <b>"A martingale characterization of mixed Poisson processes"</b>			
12. PERSONAL AUTHOR(S) <b>Dietmar Pfeifer</b>			
13a. TYPE OF REPORT <b>technical</b>	13b. TIME COVERED FROM <b>9/1/85</b> TO <b>8/31/86</b>	14. DATE OF REPORT (Yr., Mo., Day) <b>October 1985</b>	15. PAGE COUNT <b>8</b>
16. SUPPLEMENTARY NOTATION <b>Mixed Poisson process, martingales, intensities</b>			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) It is shown that an elementary pure birth process is a mixed Poisson process iff the sequence of post-jump intensities forms a martingale with respect to the $\sigma$ -fields generated by the jump times of the process. In this case, the post-jump intensities converge a.s. to the mixing random variable of the process.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>	
22a. NAME OF RESPONSIBLE INDIVIDUAL <b>Major Woodruff</b>		22b. TELEPHONE NUMBER (Include Area Code) <b>703-257-5027</b>	22c. OFFICE SYMBOL <b>7171</b>

MIC FILE COPY

## CENTER FOR STOCHASTIC PROCESSES

Department of Statistics  
University of North Carolina  
Chapel Hill, North Carolina



A martingale characterization of mixed Poisson processes

by

Dietmar Pfeifer

Technical Report No. 119

October 1985

85 12 31 065

-A-

# A MARTINGALE CHARACTERIZATION OF MIXED POISSON PROCESSES

by

Dietmar Pfeifer\*

Technical University Aachen

and

University of North Carolina at Chapel Hill

and

Ursula Heller

Technical University Aachen

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	



## Abstract

It is shown that an elementary pure birth process is a mixed Poisson process <sup>if</sup> ~~the~~ the sequence of post-jump intensities forms a martingale with respect to the <sup>delta</sup> ~~g~~-fields generated by the jump times of the process. In this case, the post-jump intensities converge a.s. to the mixing random variable of the process. *Keywords: applied probability.*

MIXED POISSON PROCESS, MARTINGALES, INTENSITIES

\*Research supported in part by US AFOSR Contract No. F 49620 85 C 0144.

AIR FORCE OFFICE OF  
NOTICE  
This  
NOTE  
Distri  
MATTHEW  
Chief, Technical Division

## 1. Introduction

Mixed Poisson processes play an important role in many branches of applied probability, for instance in insurance mathematics and physics (see Albrecht (1985) and Pfeifer (1986) for recent surveys). They belong to the class of elementary pure birth processes  $\{N(t); t \geq 0\}$  with standard transition probabilities

$$p_{nm}(s, t) = P(N(t) = m \mid N(s) = n), \quad 0 \leq n \leq m, \quad 0 \leq s \leq t, \quad (1)$$

possessing right-continuous paths and positive and finite birth rates

$$\lambda_n(t) = \lim_{h \downarrow 0} \frac{1}{h} p_{n, n+1}(t, t+h), \quad n, t \geq 0, \quad (2)$$

and all finite-dimensional marginals of the jump-time sequence  $\{T_n; n \geq 0\}$  are absolutely continuous with respect to Lebesgue measure (see Pfeifer (1982)). For such processes, the jump times form a Markov chain with transition probabilities

$$P(T_n > t \mid T_{n-1} = s) = \frac{1 - F_n(t)}{1 - F_n(s)}, \quad 0 \leq s \leq t, \quad n \geq 1 \quad (3)$$

and initial distribution function  $F_0$  where

$$1 - F_n(x) = \exp \left[ - \int_0^x \lambda_n(u) du \right], \quad x \geq 0, \quad n \geq 0 \quad (4)$$

(the  $F_n$  are in fact all cumulative distribution functions), hence all  $F_n$

are absolutely continuous with densities  $f_n$  (say), and the conditional densities for the transition probabilities can be represented as

$$f_n(s | t) = \frac{f_n(s)}{1 - F_n(t)}, \quad 0 \leq t \leq s, n \geq 1. \quad (5)$$

Moreover, the birth rates coincide with the hazard rates

$$\lambda_n(t) = \frac{f_n(t)}{1 - F_n(t)} \quad \text{a.e.,} \quad n, t \geq 0. \quad (6)$$

If especially  $\{N(t); t \geq 0\}$  is a mixed Poisson process, then also

$$\lambda_n(t) = \frac{\int_0^\infty x^{n+1} e^{-xt} dG(x)}{\int_0^\infty x^n e^{-xt} dG(x)}, \quad n, t \geq 0, \quad (7)$$

where  $G$  is the cdf of the mixing random variable  $\Lambda$  (say). In fact, Lundberg (1940) has proved that such a representation characterizes the intensities of a mixed Poisson process.

In terms of random variables, a mixed Poisson process behaves like a homogeneous Poisson process with rate  $\lambda$  given  $\Lambda = \lambda$ , from which it also follows that

$$\lambda_n(t) = E(\Lambda | N(t) = n), \quad n, t \geq 0. \quad (8)$$

The following result completes some of Lundberg's (1940) results on the asymptotic behaviour of the intensities for mixed Poisson processes.

Lemma. Let  $\{t_n; n \geq 1\}$  be a sequence of positive real numbers converging to  $t > 0$  such that  $|\frac{t_n}{t} - 1| = o(n^{-1/3}), n \rightarrow \infty$ . Then, if  $1/t$  is a point of increase of  $G$ , we have

$$\lim_{n \rightarrow \infty} \lambda_n(nt_n) = \frac{1}{t}. \quad (9)$$

Proof. Let  $\epsilon_n > 0$  be chosen in such a way that  $n \epsilon_n^3 \rightarrow 0, n \epsilon_n^2 \rightarrow \infty$  and  $(\frac{t_n}{t} - 1)/\epsilon_n \rightarrow 0$  for  $n \rightarrow \infty$ . From relation (7) it follows that

$$\begin{aligned} \lambda_n(nt_n) &= \frac{1}{t_n} \int_0^\infty e^{n(xt_n)^{n+1}} e^{-nxt_n} dG(x) \bigg/ \int_0^\infty e^{n(xt_n)^n} e^{-nxt_n} dG(x) \\ &\sim \frac{1}{t_n} \int_{(1-\epsilon_n)/t_n}^{(1+\epsilon_n)/t_n} xt_n \exp\left(-\frac{n}{2}(1-xt_n)^2\right) dG(x) \bigg/ \int_{(1-\epsilon_n)/t_n}^{(1+\epsilon_n)/t_n} \exp\left(-\frac{n}{2}(1-xt_n)^2\right) dG(x) \\ &\sim \frac{1}{t_n} \quad \text{for } n \rightarrow \infty. \quad \text{This proves the Lemma.} \end{aligned}$$

The above result is in general not true without further conditions on  $G$  as can e.g. be seen by mixing distributions concentrated in a single point  $\lambda > 0$ ; here  $\lambda_n(t) = \lambda$  for all  $n$  and  $t$ ,  $\lambda$  being the only point of increase of  $G$ . As an example, consider a Pólya-Lundberg process where  $\Lambda$  follows a gamma distribution with mean  $\mu > 0$  and variance  $\alpha \mu^2$ ,  $\alpha > 0$ . Here

$$\lambda_n(t) = \mu \frac{1 + \alpha_n}{1 + \mu \alpha t}, \quad n, t \geq 0, \quad (10)$$

from which the validity of (9) can be seen explicitly, for all  $t > 0$ .



## 2. The martingale characterization

Let  $\{\lambda_n(T_{n-1}); n \geq 1\}$  denote the sequence of post-jump intensities.

In the light of (2), the post-jump intensities describe the transition behaviour of the process immediately after a jump has occurred. The following result gives a characterization of mixed Poisson processes by a martingale property of this sequence.

Theorem 1 Let  $\{N(t); t \geq 0\}$  be an elementary pure birth process with intensities  $\{\lambda_n(t); n, t \geq 0\}$  and jump times  $\{T_n; n \geq 0\}$ . Let for  $n \geq 1$  denote  $A_n$  the  $\sigma$ -field generated by  $T_0, \dots, T_{n-1}$ . Then  $\{N(t); t \geq 0\}$  is a mixed Poisson process iff the post-jump intensities  $\{\lambda_n(T_{n-1}); n \geq 1\}$  form a martingale with respect to  $\{A_n; n \geq 1\}$ .

Proof. Due to the Markov structure of jump times the martingale property of the post-jump intensities is equivalent to

$$E(\lambda_{n+1}(T_n) \mid T_{n-1} = t) = \lambda_n(t) \quad \text{a.s. for all } n \geq 1 \quad (11)$$

which by (5) and (6) is in turn equivalent to

$$\int_t^\infty \frac{f_{n+1}(s)}{1-F_{n+1}(s)} \frac{f_n(s)}{1-F_n(s)} ds = \frac{f_n(t)}{1-F_n(t)} \quad \text{a.e.} \quad (12)$$

saying that the density  $f_n$  is differentiable a.e. with

$$f'_n(t) = - \frac{f_{n+1}(t)}{1-F_{n+1}(t)} f_n(t) \quad \text{a.e.} \quad (13)$$

or equivalently

$$\frac{d}{dt} \log(1-F_{n+1}(t)) = - \frac{f_{n+1}(t)}{1-F_{n+1}(t)} = \frac{f'_n(t)}{f_n(t)} = \frac{d}{dt} \log f_n(t) \quad \text{a.e.} \quad (14)$$

Integration of this last relation shows that there are constants  $c_n > 0$  such that

$$1-F_{n+1}(t) = c_n f_n(t), \quad t \geq 0, \quad (15)$$

which in turn implies that  $f_n$  is absolutely continuous and the recursive formula

$$f_{n+1}(t) = -c_n f'_n(t) \quad (16)$$

holds everywhere on  $[0, \infty)$ . By induction, we see that all derivatives of  $f_n$  exist on  $[0, \infty)$ , and that for all  $n \geq 1$ ,

$$f_n(t) = (-1)^n \prod_{k=0}^{n-1} c_k f_0^{(n)}(t), \quad t \geq 0. \quad (17)$$

Since by assumption, the intensities (and hence all  $f_n$ ) are positive and finite, we have

$$(-1)^n f_0^{(n)}(t) \geq 0, \quad n, t \geq 0. \quad (18)$$

The density  $f_0$  thus is completely monotonic on  $[0, \infty)$ , hence by Bernstein's (1928) theorem there is a bounded and non-decreasing right-continuous function  $H$  such that

$$f_0(t) = \int_0^\infty e^{-xt} dH(x), \quad t \geq 0. \quad (19)$$

In fact, since  $f_0$  is a density, we have that  $\frac{1}{x} dH(x) = dG(x)$  is a probability measure from which it follows that

$$\begin{aligned} f_n(t) &= \prod_{k=0}^{n-1} c_k \int_0^{\infty} x^{n+1} e^{-xt} dG(x) \\ 1-F_n(t) &= \prod_{k=0}^{n-1} c_k \int_0^{\infty} x^n e^{-xt} dG(x), \quad n, t \geq 0. \end{aligned} \quad (20)$$

Hence relation (7) is satisfied, saying that  $\{N(t); t \geq 0\}$  must be a mixed Poisson process with mixing distribution  $dG(x)$ .

Conversely, since every mixed Poisson process has intensities of the form (7), it is easily seen that relation (12) holds, hence the post-jump intensities possess the martingale property, which proves the theorem.

It should be pointed out that since  $f_0(0) < \infty$  by our assumptions, the mixing random variable must be integrable with

$$E(\Lambda) = f_0(0). \quad (21)$$

A simple application of the Martingale Convergence Theorem (see e.g. Billingsley (1979)) then shows that the post-jump intensities converge a.s. to some integrable random variable since also

$$E(\lambda_1(T_0)) = \int_0^{\infty} \frac{f_1(t) f_0(t)}{1-F_1(t)} dt = \int_0^{\infty} -\frac{f_0'(t)}{f_0(t)} f_0(t) dt = f_0(0). \quad (22)$$

The question now is what the possible limits of the post-jump intensities are. The following result gives an answer to this.

Theorem 2. If  $\Lambda$  is the mixing random variable of the process, then the post-jump intensities converge a.s. to  $\Lambda$ .

Proof. For any mixed Poisson process, we have  $(n+1)/T_n \rightarrow \Lambda$  a.s. by the strong law of large numbers, applied to the Poisson process with rate  $\lambda$ , conditionally on  $\Lambda = \lambda$ , and by the law of the iterated logarithm,

$|\lambda \frac{T_n}{n+1} - 1| = O(\sqrt{\frac{1}{n} \log \log n}) = o(n^{-1/3})$  a.s. for  $n \rightarrow \infty$ . Since also  $\Lambda$  is a.s. concentrated on the points of increase of  $G$ , the cdf of  $\Lambda$ , we have by the above Lemma

$$\lambda_{n+1}(T_n) = \lambda_{n+1}((n+1)(T_n/n+1)) \sim (n+1)/T_n \rightarrow \Lambda \quad \text{a.s.}, \quad (23)$$

which proves the theorem.

For instance, if  $\Lambda$  is concentrated on two points  $v_1 < v_2$  with mass  $\alpha$  and  $1-\alpha$  each ( $\alpha > 0$ ), then

$$\lambda_n(nt) \rightarrow \begin{cases} v_2, & t \leq l(v_1, v_2) \\ \alpha v_1 + (1-\alpha)v_2, & t = l(v_1, v_2) \\ v_1, & t \geq l(v_1, v_2) \end{cases} \quad (24)$$

where  $l(v_1, v_2) = (\log v_2 - \log v_1)/(v_2 - v_1)$ , as can be seen from Lundberg (1940), relation (108). Since always  $1/v_2 < l(v_1, v_2) < 1/v_1$ , it can explicitly be seen that

$$\lambda_{n+1}(T_n) \rightarrow \begin{cases} v_1 & \text{with probability } \alpha \\ v_2 & \text{with probability } 1-\alpha, \end{cases} \quad (25)$$

i.e.  $\lambda_{n+1}(T_n) \rightarrow \Lambda$  a.s.

Acknowledgements. We would like to thank P. Deheuvels for some stimulating suggestions concerning an earlier draft of the paper.

References.

Albrecht, P. (1985). Mixed Poisson processes. In: Encyclopedia of Statistical Sciences, Vol. 6, Wiley, N.Y.

Bernstein, S. (1928). Sur les fonctions absolument monotones. Acta Mathematica 51, 1 - 66.

Billingsley, P. (1979). Probability and Measure. Wiley, N.Y.

Lundberg, O. (1940). On Random Processes and Their Application to Sickness and Accident Statistics. Reprinted by Almqvist and Wiksells (1964), Uppsala.

Pfeifer, D. (1982). The structure of elementary pure birth processes. J.Appl. Prob.19, 664- 667.

Pfeifer, D. (1986). Pólya-Lundberg process. To appear in: Encyclopedia of Statistical Sciences, Vol. 7, Wiley, N.Y.

**END**

**FILMED**

---

*2-86*

**DTIC**